

Information For the User
Flavoxate (200 mg) + nitrofurantoin (100 mg) Tablet

Uriviga N Tablet

Uriviga N Tablet

Dosage Form: Tablet

Composition:

Each film-coated tablet contains:

- Flavoxate Hydrochloride IP 200 mg
- Nitrofurantoin IP 100 mg

Description

Uriviga N Tablet is a fixed-dose combination of Flavoxate Hydrochloride, a urinary antispasmodic agent, and Nitrofurantoin, a urinary antibacterial. It is indicated for the management of urinary tract infections (UTIs) associated with urinary discomfort. The combination provides symptomatic relief from urinary spasms while effectively treating bacterial infection of the urinary tract.

Indications

Indicated for:

- Acute and chronic urinary tract infections
- Dysuria, urinary frequency, and urgency associated with UTIs
- Cystitis and other lower urinary tract infections

It helps in relieving bladder spasms, reducing pain during urination, and eliminating susceptible urinary pathogens.

Dosage & Administration

- Dosage should be as directed by the physician.
- Swallow the tablet whole with water, preferably after meals.
- Complete the full course of antibiotic therapy as prescribed.

Pharmacological Action

- **Flavoxate Hydrochloride:** Acts directly on smooth muscle of the urinary tract, relieving spasms and reducing urinary urgency and discomfort.
- **Nitrofurantoin:** Inhibits bacterial enzyme systems and interferes with bacterial protein synthesis, leading to bactericidal activity against susceptible urinary pathogens.

The combination provides both antimicrobial action and symptomatic relief.

Side Effects

Common side effects may include:

- Nausea and vomiting
- Abdominal discomfort
- Headache
- Dizziness
- Allergic reactions (rare)

Precautions & Warnings

- Contraindicated in patients with significant renal impairment
- Use with caution in elderly patients
- Not recommended during late pregnancy
- Monitor for signs of pulmonary or hepatic reactions with prolonged use
- Use under strict medical supervision

Drug Interactions

- Antacids containing magnesium may reduce absorption of Nitrofurantoin
- Caution with other drugs affecting renal function
- Consult physician before co-administration with other urinary agents

Storage

- Store below 25°C in a cool, dry place
- Protect from light and moisture
- Keep out of reach of children

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for