

Information For the User
Ursodeoxycholic acid 300 mg syrup

Uriviga 300 Syrup

Brand Name: Uriviga 300 Syrup

Dosage Form: Syrup

Therapeutic Category: Hepatoprotective / Bile Acid – Gallstone Dissolution Agent

Composition:

Each 5 ml contains:

- **Ursodeoxycholic Acid IP** – 300 mg
- Excipients – q.s.
- Colour: Approved colour used
- Flavour: Palatable base

Pharmacology & Mechanism of Action:

Ursodeoxycholic acid (UDCA) is a naturally occurring hydrophilic bile acid. It works by reducing the cholesterol content of bile and bile stones through suppression of hepatic cholesterol synthesis and inhibition of intestinal absorption of cholesterol. It stabilizes hepatocyte canalicular membranes, stimulates bile secretion, and protects liver cells from toxic bile acids. It is effective in dissolving cholesterol gallstones and improving liver function in cholestatic conditions.

Indications:

- Cholesterol-rich gallstone dissolution
- Primary biliary cholangitis (PBC)
- Hepatobiliary disorders in chronic liver diseases
- Non-alcoholic fatty liver disease (NAFLD)
- Bile reflux gastritis

Dosage & Administration:

- **Adults:** 5–10 ml once or twice daily after meals, or as directed by the physician
- Dosage may be adjusted based on body weight and clinical condition
- Shake well before use
- To be administered under the supervision of a registered medical practitioner

Contraindications:

- Hypersensitivity to UDCA or any excipients
- Calcified or pigment gallstones
- Obstructive biliary tract diseases
- Non-functioning gallbladder
- Acute inflammation of gallbladder or bile ducts

Warnings & Precautions:

- Periodic monitoring of liver function tests is advised during therapy
- Not recommended during pregnancy unless clearly necessary
- Use with caution in lactating women
- Regular ultrasound checkups may be needed to monitor gallstone dissolution

Drug Interactions:

- Cholestyramine, colestipol, or antacids containing aluminium may reduce UDCA absorption – administer at least 2 hours apart
- May alter absorption of ciclosporin and antibiotics – monitor levels accordingly

Adverse Effects:

- Occasional: Diarrhea, nausea, mild abdominal pain
- Rare: Calcification of gallstones, skin rash, pruritus
- Long-term therapy may require liver enzyme monitoring

Storage:

- Store below 25°C
- Protect from light and moisture
- Keep out of reach of children

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for