

Virafly D tablet

INFORMATION FOR THE USE

Virafly D Tablet

Sildenafil Citrate 50mg + Dapoxetine 30mg – Tablet

Description:

Virafly D Tablet combines Sildenafil Citrate and Dapoxetine, offering dual action for the management of male sexual disorders. Sildenafil enhances erectile function by increasing blood flow to the penis, while Dapoxetine is a short-acting selective serotonin reuptake inhibitor (SSRI) used to treat premature ejaculation. This synergistic combination addresses both erectile dysfunction (ED) and premature ejaculation (PE) in a single formulation.

Composition:

- Sildenafil Citrate: 50mg
- Dapoxetine Hydrochloride: 30mg

Indications:

1. Erectile Dysfunction (ED) – for improving the ability to achieve and maintain an erection
2. Premature Ejaculation (PE) – for delaying ejaculation and improving control
3. Combined management of ED and PE in adult males
4. Enhancing sexual confidence and performance

Mechanism of Action:

- **Sildenafil Citrate:** A PDE5 inhibitor that increases cGMP levels by inhibiting its degradation. This leads to smooth muscle relaxation and increased blood flow to the penile tissues during sexual stimulation, enabling erection.
- **Dapoxetine:** Inhibits serotonin reuptake at synapses, prolonging ejaculation latency time, enhancing control over ejaculation, and reducing distress associated with PE.

Dosage and Administration:

- Take one tablet orally with a full glass of water, 1 to 3 hours before anticipated sexual activity.
- Do not exceed one dose in 24 hours.
- Should be taken on an empty stomach for optimal effect.
- Sexual stimulation is required for efficacy.

Side Effects:

- Common: Headache, nausea, dizziness, nasal congestion, flushing, indigestion
- Less Common: Blurred vision, anxiety, insomnia
- Rare but Serious: Priapism (prolonged erection), severe hypotension, serotonin syndrome (with Dapoxetine)

Precautions:

1. Not for use in individuals below 18 years or above 65 without medical supervision
2. Avoid concurrent use with nitrates or nitric oxide donors (risk of severe hypotension)
3. Not recommended in patients with significant cardiovascular disorders, liver/kidney impairment
4. Avoid alcohol and recreational drugs as they may enhance side effects
5. Monitor for mood changes or depressive symptoms due to SSRI component

Storage:

- Store in a cool, dry place below 25°C, away from moisture and direct sunlight
- Keep out of reach of children

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for