Information For the Use Methylprednisolone (16mg)

Volipred 16 Tablet

Information for the Use

Methylprednisolone (16 mg) Tablet

• Other Conditions: Certain cancers, adrenal insufficiency, and organ transplant rejection prophylaxis.

Composition:

Each tablet contains:

Methylprednisolone: 16 mg

Description:

Methylprednisolone is a synthetic glucocorticoid with potent anti-inflammatory and immunosuppressive properties. It works by modulating the immune response and reducing inflammation, making it effective in treating a wide range of conditions.

Indications:

Methylprednisolone 16 mg tablets are indicated for:

- Inflammatory Disorders: Rheumatoid arthritis, osteoarthritis, and ankylosing spondylitis.
- Autoimmune Diseases: Systemic lupus erythematosus (SLE), dermatomyositis, and vasculitis.
- Allergic Conditions: Severe asthma, allergic rhinitis, and anaphylaxis.
- Skin Disorders: Psoriasis, eczema, and dermatitis.
- Respiratory Disorders: Chronic obstructive pulmonary disease (COPD) exacerbations.
- Gastrointestinal Conditions: Crohn's disease and ulcerative colitis.
- Neurological Conditions: Multiple sclerosis exacerbations.

Mechanism of Action:

Methylprednisolone acts by binding to glucocorticoid receptors, which modulate gene expression to suppress the immune response and reduce the production of inflammatory mediators such as prostaglandins and cytokines.

Dosage and Administration:

- General Dosage: Dosage varies depending on the condition being treated, severity, and patient response.
- Recommended Dosage for Adults:
 - Initial dose: 16 mg to 48 mg per day, divided into 1 to 4 doses.
 - Maintenance dose: Gradually reduce to the lowest effective dose.
- Pediatric Use: Not recommended without specific medical advice.
- Take the tablet orally with food to reduce gastrointestinal upset.

Contraindications:

- Hypersensitivity to Methylprednisolone or its components.
- Systemic fungal infections.
- Live or attenuated vaccines during treatment.

Precautions:

- Infections: May mask symptoms of infection; use cautiously in patients with active infections.
- Bone Health: Long-term use may lead to osteoporosis.
- Blood Sugar: Monitor in diabetic patients as it may increase blood glucose levels.
- Eye Conditions: Use cautiously in patients with glaucoma or cataracts.
- Psychiatric Effects: Monitor for mood swings, depression, or psychosis in susceptible patients.
- Pregnancy and Lactation:
 - Use only if the benefits outweigh the risks during pregnancy.
 - Small amounts may pass into breast milk; use cautiously while breastfeeding.

Possible Side Effects: Common side effects:

- Nausea, vomiting, or abdominal discomfort.
- Weight gain, fluid retention, or swelling.
- Mood changes, irritability, or insomnia.

Serious side effects:

- Infections: Increased susceptibility to infections.
- Adrenal Suppression: Risk of adrenal insufficiency with sudden discontinuation.
- Osteoporosis: Long-term use may weaken bones.
- Gastrointestinal Complications: Peptic ulcers or bleeding.

Drug Interactions:

NSAIDs: Increases the risk of gastrointestinal bleeding.

- Antidiabetic Agents: May reduce their effectiveness, requiring dosage adjustments.
- Live Vaccines: Should be avoided during treatment.
- Cyclosporine and Warfarin: Monitor for altered effects.

Storage:

- Store at 15°C to 25°C.
- Protect from light and moisture.
- · Keep out of reach of children.

Special Instructions:

- Do not stop the medication abruptly; taper the dose under medical supervision.
- Regular monitoring of blood pressure, blood glucose, and bone density is recommended for long-term users.
- Take the medication as directed, and do not exceed the prescribed dose.

Note: Always consult your healthcare provider for specific advice regarding the use of Methylprednisolone 16 mg tablets. This medication should be used under strict medical supervision.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for