

Fungal Diastase (1:800) 100mg + Papain 60mg + Activated Charcoal 75mg

Zymvigo Tablet

Fungal Diastase (1:800) 100mg + Papain 60mg + Activated Charcoal 75mg Tablet Information

Generic Name:

Fungal Diastase (1:800) + Papain + Activated Charcoal

Strength:

Fungal Diastase 100mg, Papain 60mg, Activated Charcoal 75mg

Form:

Tablet

Therapeutic Class:

Digestive Enzyme Supplement, Antacid, and Adsorbent

Description

This combination tablet contains:

- **Fungal Diastase:** A fungal-derived enzyme that breaks down starches into simpler sugars, aiding in the digestion of carbohydrates.
- **Papain:** A proteolytic enzyme from papaya that helps in the breakdown of proteins, promoting overall digestion and reducing bloating or indigestion.
- **Activated Charcoal:** A natural adsorbent that binds to toxins, gases, and other unwanted substances in the stomach and intestines, helping to alleviate bloating, gas, and discomfort.

Together, these ingredients support digestive health by improving enzyme activity and reducing bloating or discomfort caused by indigestion or gas.

Indications

Fungal Diastase + Papain + Activated Charcoal 100mg + 60mg + 75mg tablets are indicated for:

- **Indigestion:** Relief of symptoms such as bloating, gas, and fullness after meals.
 - **Post-meal discomfort:** Help in the digestion of food, particularly after heavy or rich meals.
 - **Flatulence:** Reduction of excessive gas formation in the stomach and intestines.
 - **Dyspepsia:** General digestive discomfort due to impaired digestion.
 - **Acid reflux:** Reducing symptoms of acid reflux by improving the digestion process and reducing gas formation.
-

Dosage and Administration

- **Adults:**
 - 1-2 tablets after meals, or as directed by a healthcare provider.
- **Children (under 12 years):**
 - Consult a healthcare professional before use.

Note: It is recommended to take the tablet with water to help with swallowing and digestion.

Contraindications

- **Hypersensitivity to any of the ingredients in the tablet.**
- **Use in patients with gastrointestinal perforation, active peptic ulcers, or severe gastrointestinal disorders should be avoided.**
- **Not recommended for use in individuals with known allergy to papaya or fungal enzymes.**

- **Activated Charcoal: May reduce the effectiveness of certain medications by absorbing them, including anticoagulants, antidepressants, and antidiabetic drugs.**
- **Enzyme Supplements: Use with caution if taking other digestive enzymes concurrently to avoid potential overdose or side effects.**

Precautions

- **Gastrointestinal Disorders: Use cautiously in patients with any significant gastrointestinal conditions, such as ulcerative colitis, Crohn's disease, or history of gastrointestinal surgery.**
- **Pregnancy and Lactation: Consult a healthcare provider before using during pregnancy or breastfeeding.**
- **Drug Interactions: Activated charcoal may interfere with the absorption of certain medications, so it should be taken at a different time from other drugs.**

Storage Instructions

- **Store in a cool, dry place at room temperature (15-25°C).**
- **Keep away from moisture and light.**
- **Keep out of reach of children.**

Side Effects

Common side effects:

- **Mild gastrointestinal disturbances such as nausea or upset stomach.**
- **Diarrhea or constipation in some cases due to the activated charcoal.**

Rare but serious side effects:

- **Allergic reactions such as rash, itching, or swelling.**
- **Severe gastrointestinal discomfort, bloating, or constipation.**

Additional Notes

- **This product is intended as a digestive aid and should not replace a balanced diet or medical treatment for gastrointestinal diseases.**
- **If symptoms persist or worsen, consult a healthcare provider.**
- **The use of activated charcoal may cause a temporary blackening of the stool.**

Always follow your healthcare provider's instructions regarding use.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for

Drug Interactions